



■ Embella



Embella[®]
(deoxycholic acid) injection

Tight and Trim, Like your Dream







Indications

FDA-approved for submental convexity associated with moderate to severe submental fat based on Clinician-Reported Submental Fat Rating Scale (CS-SMFRS):

Scale	0	1	2	3	4
Submental Convexity	Absent	Mild	Moderate	Severe	Extreme
Description	No localized submental fat evident	Minimal localized submental fat	Prominent localized submental fat	Marked localized submental fat	Extreme submental convexity
Representative Photographs					

Limitations

Do not inject above the inferior border of the mandible.
Do not inject within a region defined by 1-1.5 cm line below the inferior border.
Avoid injection into the platysma






Indications

FDA-approved for submental convexity associated with moderate to severe submental fat
Clinician-Reported Submental Fat Rating Scale (CS-SMFRS):

Scale	0	1	2	3	4
Submental Convexity	Absent	Mild	Moderate	Severe	Extreme
Description	No localized submental fat evident	Minimal localized submental fat	Prominent localized submental fat	Marked localized submental fat	Extreme submental convexity
Representative Photographs					

Limitations

Do not inject above the inferior border of the mandible.
Do not inject within a region defined by 1-1.5 cm line below the inferior border.
Avoid injection into the platysma





- Lipolytic agent
- 10 mg/ml Deoxycholic acid + 6.4mg/ml HA
- Subcutaneous use **ONLY**
- 2-4 treatment sessions(maximum:6 treatments) with 4 weeks intervals
- Apply a 1 cm injection grid to mark the injection sites.
Inject 0.2 ml (2mg) per injection site. (maximum: 10 ml=100 mg= 50 injections)

ADVERSE REACTIONS:

- Very common ($\geq 10\%$):
- Pruritus
- Paresthesia
- Adverse reactions at injection site, such as:
 - Bruising
 - erythema
 - hematoma
 - induration
 - nodule
 - numbness
 - swelling

Precautions:

Patients with excessive skin laxity, Prominent platysma bands

